WORLD INTELLECTUAL PROPERTY ORGANIZATION



B2

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 4:		(11) International Publication Number: WO 85/ 04100
A61K 9/40, 9/22, 9/52	A1	(43) International Publication Date: 26 September 1985 (26.09.85)
(21) International Application Number: PCT/US (22) International Filing Date: 22 February 1985 ((31) Priority Application Number: (32) Priority Date: 21 March 1984 (591,7 (21.03.8	New York, NY 10017 (US). (81) Designated States: AU, DK, FI, JP, KR.
 (71) Applicant: AMERICAN HOME PRODUCT PORATION [US/US]; 685 Third Avenue, Now 10017 (US). (72) Inventors: KALIDINDI, Sanyasi, Raju; 36B A Square Apartments, 1148 Mulberry Lane, Gone 27834 (US). GIANNINI, Robert, Pet Meadowbrook Road, East Norriton, PA 194 GASKILL, James, Lawrence; Apartment M gartown and Avon Roads, Devon, PA 193 RUSSO, Emanuel, Joseph; 8 Morris Circle PA 19087 (US). 	Arlingt reenviler; 5401 (U307, S.33)	R- k, n e, 8 8).
(54) Title: SUSTAINED RELEASE PHARMACE	UTIC	L CAPSULES
(57) Abstract		

A gastrointestinal, pH-independent, sustained-release pharmaceutical unit dosage form comprising a non-compressed mixture of a therapeutic agent and from about 10 to about 60 percent by weight of a high molecular weight hydroxypropylmethylcellulose, and a method for production thereof in hard gelatin capsules.

Ţ

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT AU BB BE BG BR CF CG CH DE DK FI FR	Austria Austrialia Barbados Belgium Bulgaria Brazil Central African Republic Congo Switzerland Cameroon Germany, Federal Republic of Denmark Finland France	GA GB HU IT JP KP KR LI LK LU MC MG ML	Gabon United Kingdom Hungary Italy Japan Democratic People's Republic of Korea Republic of Korea Liechtenstein Sri Lanka Luxembourg Monaco Madagascar Mali	MR MW NL NO RO SD SE SN SU TD TG US	Mauritania Malawi Netherlands Norway Romania Sudan Sweden Senegal Soviet Union Chad Togo United States of America

-1-

SUSTAINED RELEASE PHARMACEUTICAL CAPSULES

American Home Products Corporation, a Corporation organised and existing under the laws of the State of Delaware, United States of America of 685 Third Avenue, New York, New York 10017, United States of America.

Ť

10

15

20

25

30

35

Background of the Invention

The convenience and advantages of administering a single dose of medication which provides prolonged or sustained-release as opposed to the administration of a number of single doses at regular intervals are indisputable. Conventionally, sustained-release is achieved by controlling dissolution and/or diffusion of the medicament from the dosage form. Several carrier materials which are employed for this purpose include waxes, fatty materials, polymers, natural, synthetic and semi-synthetic gums, etc. Among the gums, hydroxypropylmethylcelluloses constitute an important class of materials because of their pH-independent effect as well as their synthetic origin as opposed to the natural gums such as alginates, Karaya, Guar, Locust bean, etc.

The use of hydroxypropylmethylcelluloses as carrier bases in sustained-action tablets has been well-established. For example, Christensen et al. (U.S. Patent 3,065,143) teach sustained-release tablets which contain a hydroxypropylmethylcellulose which, upon contact with gastric fluid, swells and forms a waterimpermeable barrier on the tablet surface thereby providing sustained release by diffusion of drug through the barrier. It is clear from their teachings that the gum has to be pressed into a tablet for it to work as a sustained-action agent. Sheth et al. (U.S. Patent 4,167,558) disclose sustained-release tablets containing hydroxypropylmethylcellulose 60 HG. Again the dosage form is tablet and in addition there is a restriction on the density of the tablets to assure buoyancy and release the whole of the medicament in the stomach. Lowey et al. (U.S. Patent 3,870,790) disclose a solid, compressed buccal product containing low molecular weight hydroxypropylmethylcellulose which has been modified by humidification and airdrying. Similarly, Schor (U.S. Patent 4,226,849) disclosed the invention of tablets, lozenges, suppositories and/or other compressed dosage forms, which have prolonged-release wherein the hydroxypropylmethylcellulose has been subjected to hydrolysis and oxidation to generate a desired minimum concentration of carbonyl and carboxyl groups. The hydroxypropylmethylcellulose used in these teachings are of low molecular weight. In another patent (U.S. Patent 4,389,393) Schor et al. disclose the use of high molecular weight hydroxypropylmethylcelluloses in low concentrations for achieving sustained drug release action from compressed solid dosage forms. The hydroxypropylmethylcelluloses used by Schor et al. have a molecular weight above 50,000, a methoxyl content of 16-24% and hydroxypropoxyl content of 4-32%. Thus, Schor et al. (4,389,393) require that the material be compressed and it is clear from these teachings and those of Christensen et al.

10

15

20

25

30

35

ţ

Ç

(U.S. Patent 3,065,143), Sheth et al. (4,167,558), Lowey et al. (3,870,790) and Schor et al. (4,226,849 and 4,389,393) that compression is essential for the formation of a diffusion barrier layer on the surface of the oral dosage form to provide sustained action.

Sheth and Tossounian (U.S. Patent 4,126,672) showed that hydroxypropyl-methylcellulose provides sustained release in a buoyant capsule dosage form. However, because of the buoyancy constraints, the density of the capsules may have to be adjusted by the use of a fatty material so that the capsules float in the gastric fluid. It is clear from their teachings that floating is essential for their concept to work. Henderson et al. (U.S. Patent 3,427,378) disclose capsules for sustained-release of medicament wherein a gum such as sodium alginate is incorporated into the pharmaceutical formulation in an amount of from 70 to 99 percent by weight and preferably 85 to 98 percent and the capsules are, as a critical feature, completely filled.

Description of the Invention

This invention provides a unit dosage form for administration of a therapeutic agent, which does not require expensive compression and granulation processing characteristic of the prior art sustained-release tablets, or density control restraints characteristic of the prior art capsules, while providing substantially the same slow drug release heretofore obtained with compressed tablets. The unit dosage form of this invention provides sustained release of drug independent of gastrointestinal pH and is therefore equally effective both in the stomach and intestine. By "independent," it is not meant that no change in dissolution rate occurs with change of pH, but that any change is so insubstantial that drug delivery is not meaningfully altered.

In essence, it has been discovered that pharmaceutical compositions formulated with relatively high molecular weight hydroxypropylmethylcellulose (>50,000) which heretofore have been employed in the production of compressed unit dosage forms such as tablets, buccal lozenges, suppositories, etc. may be used without compression in hard gelatin capsules without loss of the sustained release effect observed in compressed tablets. Although the pharmaceutical literature has well documented the necessity for compressing hydroxypropylmethylcellulose-containing drug formulations to provide an external surface barrier to drug release, or for control of the density of an encapsulated drug formulation compounded with hydroxypropylmethylcellulose to provide buoyancy and hold the capsule in the

10

15

20

25

30

35

stomach, applicants have found that neither of these preparative restrictions are necessary to provide a sustained release unit dosage capsule for administration. The formulations employed in this invention are not tailored to ensure buoyancy in gastric fluids and they are equally effective when administered in such manner as to be retained with a food bolus passing into the small intestine. No special attention or limitation as to internal porosity is required with the formulations of this invention.

Thus, in accordance with this invention, there is provided a gastrointestinal, pH-independent, sustained-release pharmaceutical unit dosage form comprising a hard gelatin capsule containing a pulverulent admixture of a therapeutically active medicament; from about 10 to about 60 percent by weight of at least one hydroxy-propylmethylcellulose possessing a methoxy content of 16 to 24 weight percent, a hydroxypropoxyl content of 4 to 32 weight percent and a number average molecular weight of at least 50,000; and a pharmaceutically acceptable, nonpharmacologically functional adjuvant therefor.

The hydroxypropylmethylcelluloses employed in production of the sustainedrelease capsules are commercially available. In essence, any hydroxypropylmethylcellulose (HPMC) possessing the characteristics of Methocel & K4M, K15M and K100M are applicable, individually and as mixtures. Each of these hydrophilic materials have a methoxy content of about 16 to about 24 weight percent, a hydroxypropyl content of about 4 to about 32 weight percent and a number average molecular weight of at least 50,000. The proportion of hydroxypropylmethylcellulose to total capsule weight may vary from about 10 to about 60 percent to provide release rates of drug over a 6 to about 12 hour period. Although the release rates achieved with hard gelatin capsules containing drug formulations and hydroxypropylmethylcellulose in accordance with this invention will differ somewhat depending upon the specific drug being compounded, the sustained release effect is readily achieved. Modification of the release rate may be attained by altering the hydroxypropylmethylcellulose to drug ratio within the limits indicated above. Generally, an increased content of hydroxypropylmethylcellulose will prolong the release rate of a given drug as may be seen from the experimental data presented, infra. An optimum release rate is obtained when near total dissolution of the therapeutic agent at a substantially uniform rate is achieved over a period of from six to twelve hours and preferably in about eight hours.

Any therapeutic agent suitable for oral administration may be formulated in conformance with this disclosure and encapsulated to provide a sustained release

10

15

20

30

7.

unit dosage capsule. Thus, tranquilizers, analgesics, antihypertensives and similar drugs may be readily formulated for use in sustained release capsule form.

In addition to the therapeutic agent and hydroxypropylmethylcellulose, other conventional additives such as fillers and binders (microcrystalline cellulose, lactose, dicalcium phosphate dihydrate, etc.); lubricants (magnesium stearate, stearic acid, etc.); glidants (colloidal silicon dioxide, talc, etc.); hydrophilic gums (methylcellulose, hydroxypropylcellulose, hydroxyethylcellulose, sodium carboxymethylcellulose, etc.); disintegrants (starch, sodium starch glycolate, etc.); preservatives (methylparahydroxy benzoate, benzoic acid, etc.); antioxidants (ascorbic acid, sodium bisulfite, etc.) and colorants (certified dyes) may be employed in the formulations for filling hard gelatin capsules without materially changing the sustained release rate of medicament achieved with the indicated high molecular weight hydroxypropylmethylcellulose material.

The pharmaceutical compositions of this invention are prepared by blending an admixture of the ingredients and filling hard gelatin capsules therewith by hand or machine. Where the particulate size of any one component of the composition, or the mixture in its entirety, is of such size as to detract from the production of a homogeneous or near homogeneous blend for subsequent processing, or the desired performance of the finished dosage form (e.g. dissolution, stability, uniformity, weight variation, etc.), that component or the entire mixture may be milled to any desired size prior to a final blending and capsule filling.

The following examples illustrate several specific formulations for filling hard gelatin capsules to produce sustained release doses of the indicated therapeutic agents. For purposes of this illustration, the drugs used were:

25 Oxazepam 7-chloro-1,3-dihydro-3-hydroxy-5-phenyl-2H-1,4-benzodiazepin-2-one;

Ciramadol 1-cis-2-(a-dimethylamino-m-hydroxybenzyl)cyclohexanol;

Guanabenz (E)-[(2,6-dichlorobenzylidene)amino lguanidine; and

Lorazepam 7-chloro-5-(o-chlorophenyl)-1,3-dihydro-3-hydroxy-2H-1,4-benzo-diazepin-2-one.

374.50

30 .

- 6 -

Example 1 Hydroxypropylmethylcellulose 10 percent

	Hydroxypropylmethylcellulose 10 percent	
		mg
	Oxazepam USP	30.00
	Hydroxypropylmethylcellulose 2208 USP	37.50
	Microcrystalline Cellulose NF	94.00
	Lactose USP Hydrous	206.75
	Colloidal Silicon Dioxide NF	3.75
	Magnesium Stearate NF	3.00
	magnesium becarato 1.1	<u></u>
•	TOTAL	375.00
	Example 2	
	Hydroxypropylmethylcellulose 20 percent	
		mg
	Oxazepam USP	30.00
	Hydroxypropylmethylcellulose 2208 USP	75.00
	Microcrystalline Cellulose NF	94.00
	•	169.25
	Lactose USP Hydrous	3.75
	Colloidal Silicon Dioxide NF	3.00
	Magnesium Stearate NF	
	TOTAL	375.00
	Example 3	
	Hydroxypropylmethylcellulose 30 percent	
		mg
	Oxazepam USP	30.00
	Hydroxypropylmethylcellulose 2208 USP	112.00
	Microcrystalline Cellulose NF	94.00
	Lactose USP Hydrous	131.75
	Colloidal Silicon Dioxide NF	3.75
	Magnesium Stearat NF	3.00

TOTAL

Example 4 Hydroxypropylmethylcellulose 25 percent

		mg
	Ciramadol Hydrochloride	68.80
5	Hydroxypropylmethylcellulose 2208 USP	. 112.50
	Microcrystalline Cellulose NF	110.26
	Lactose USP Hydrous	152.27
	Colloidal Silicon Dioxide NF	4.50
	Magnesium Stearate NF	1.67
10	TOTAL	450.00
	Example 5	
	Hydroxypropylmethylcellulose 60 percent	ma .
	•	mg 20.20
	Guanabenz Acetate	90.00
15	Hydroxypropylmethylcellulose 2208 USP	39.05
	Lactose USP Hydrous	0.75
	Magnesium Stearate NF	
	TOTAL	150.00
	Example 6	
20	Hydroxypropylmethylcellulose 40 percent	
		mg
	Lorazepam	2.00 40.00
	Hydroxypropylmethylcellulose 2208 USP	
	Microcrystalline Cellulose NF	30.00
25	Lactose USP Hydrous	27.50
	Magnesium Stearate NF	0.50
	TOTAL	100.00

10

15

25

In each of the preceding examples, all the ingredients were mixed together in a Twin-shell blender for 10 minutes, milled through a Fitz Mill[®], and mixed again for 10 minutes in the same blender. The blend was then filled into hard gelatin capsules using a Zanasi LZ64 capsule-filling machine.

The above formulations provide sustained-release for 6-12 hours in vitro as shown in Table I. The dissolution rate study employed to obtain the data reported in Table I was performed in an apparatus conforming to that described as Apparatus 2 in U.S. Pharmacopeia XX page 959 (1980) by the method therein described, at 50 r.p.m. or 100 r.p.m. The dissolution medium was 500 ml 0.1N HCl for the first hour after which 500 ml phosphate buffer was added to afford a pH of 7.4, comparable to that found in the intestine. A spring coil was employed to sink the capsules and insure that they did not float.

Table I: Dissolution Data for Various Sustained-Action Capsules
Containing Hydroxypropylmethylcellulose Carrier Base.
(% Drug Dissolved as an Average of Six Capsules)

	Formulation of			•			•
-	Example No.	I Hour	2 Hours	4 Hours	6 Hours	8 Hours	12 Hours
		20	57	90	97	_	-
	2	26	56	. 79	86	88	94
20	3	15	36	55	69	79	92
•	4	37	60	. 82	95	100	-
	5	28	47	68	84	93	94
	6	20	32	50	67	82	94

To illustrate the comparable dissolution rates between compressed tablets and the capsules of this invention employing the same formulation, the following data was obtained with the appropriate formulations disclosed, <u>supra</u>, and with that of Example 7.

Example 7
Hydroxypropylmethylcellulose 38.6 percent

	Guanabenz Acetate	20.2 mg
	Hydroxypropylmethylcellulose 2208, USP	61.7 mg
5	Magnesium stearate, NF	0.55 mg
•	Microcrystalline cellulose, NF	22.0 mg
	Lactose, USP	45.6 mg
	Sodium starch glycolate, NF	10.0 mg
		
	TOTAL	160.0 mg

The tablets were conventionally prepared and the capsules were prepared by machine-filling as described, <u>supra</u>. The procedure followed in obtaining this dissolution rate data was the same as that disclosed, <u>supra</u>, except that the study employing guanabenz were run in all acid (0.1N HCl) without addition of buffer, and all samples were run at a paddle speed of 50 RPM.

Dissolution Data Comparing Tablets and Capsules Prepared With the Same Powder Blend or Granulation. (% Dissolved as an Average of Six Capsules and Tablets)

		Oxazepam with 20% HPMC 2208 USP		Guanabenz with 38.6% HPMC 2208 USP		Ciramadol with 25% HPMC 2208 USP	
20	Time (Hours)	Ţablets	Capsules	Tablets	Capsules	Tablets	Capsules
	1	12.5	13.5	35.0	37.5	26.5	37
	2	22.5	22.5	52.0	56.0	41.5	60
	4	33.0	32.0	77.0	80.0	57.5	82
25	6	40.0	39.0	93.0	92.0	75.0	95
	8	46.0	46.0	93.0	98.0	85.0	complete
	10	52.5	52.0	95.5	98.0	93.5	complete
	12	56.5	55.5	97.0	96.0	96.5	complete

20

25

To demonstrate that there is no property imparted to the capsules of this invention by the encapsulating machine, the data presented in Table III compares the dissolution rate of hand-filled capsules with those containing the same formulations when filled with a Zanasi LZ64 encapsulating machine. The formulations employed were those shown in Examples 2 and 6.

Table III: Dissolution of Hand-Filled and Machine-Filled Capsules (% Dissolved as an Average of Six Capsules)

		Oxazepan	n (Example 2)	Lorazepan	(Example 6)
10	Time (Hours)	Hand filled	Machine filled	Hand filled	Machine filled
	1	17	26	24	20
	2	. 44	56	37	32
	4	68	79	~: 55	50
	6	. 80	86	71	67
15	8	81	88	88	82
	. 12	84	94	99	94
	20	91	98	100	94

From these data, the preferred quantity of defined hydroxypropylmethyl-cellulose for general incorporation into pharmaceutical formulations for encapsulation can be seen to be from about 10 to about 50 weight percent. For the individual drugs herein exemplified the preferred quantity of defined hydroxy-propylmethylcellulose is from about 15 to about 40 weight percent with oxazepam; about 20 to about 50 weight percent with lorazepam and from about 15 to about 50 weight percent with ciramadol.

Thus, the advantages of capsule dosage forms with no loss of sustained drugrelease rate presently found in compressed tablets, clearly characterizes the benefits of this invention.

15

20

25

30

Ĵ.

CLAIMS

- 1. A gastrointestinal pH independent sustained release pharmaceutical unit dosage form comprising a hard gelatin capsule containing a pulverulent admixture of a therapeutically active medicament; from about 10 to about 60 percent by weight of at least one hydroxypropylmethylcellulose possessing a methoxy content of 16 to 24 weight percent, a hydroxypropoxyl content of 4 to 32 weight percent and a number average molecular weight of at least 50,000; and a pharmaceutically acceptable, nonpharmacologically functional adjuvant therefor.
- A pharmaceutical unit dosage form of Claim 1 in which said hydroxy propylmethylcellulose is present in an amount of from about 10 to about 50 percent by weight.
 - 3. A sustained-release pharmaceutical composition in capsule form as claimed in Claim 1 for oral administration of oxazepam, which comprises a pharmaceutically effective amount of oxazepam admixed with a hydroxypropylmethylcellulose wherein said hydroxypropylmethylcellulose has a methoxyl content of about 16-24 percent, a hydroxypropoxyl content of about 4-32 percent and an average number molecular weight above 50,000 and said hydroxypropylmethylcellulose represents between about 10 to about 60 weight percent of the contents of said capsule.
 - 4. A pharmaceutical composition in capsule form as claimed in Claim 3 in which said hydroxypropylmethylcellulose represents between about 15 to about 40 percent of the contents of said capsule.
 - 5. A sustained-release pharmaceutical composition in capsule form as claimed in Claim 1 for oral administration of lorazepam, which comprises a pharmaceutically effective amount of lorazepam admixed with a hydroxypropyl-methylcellulose wherein said hydroxypropylmethylcellulose has a methoxyl content of about 16-24 percent, a hydroxypropoxyl content of about 4-32 percent and an average number molecular weight above 50,000 and said hydroxypropylmethyl-cellulose represents between about 10 to about 60 weight percent of the contents of said capsule.

10

- 6. A pharmaceutical composition in capsule form as claimed in Claim 5 in which said hydroxypropylmethylcellulose represents between about 20 to about 50 percent of the contents of said capsule.
- 7. A sustained-release pharmaceutical composition in capsule form as claimed in Claim 1 for oral administration of ciramadol, which comprises a pharmaceutically effective amount of ciramadol admixed with a hydroxypropylmethylcellulose wherein said hydroxypropylmethylcellulose has a methoxyl content of about 16-24 percent, a hydroxypropoxyl content of about 4-32 percent and an average number molecular weight above 50,000 and said hydroxypropylmethylcellulose represents between about 10 to about 60 weight percent of the contents of said capsule.
- 8. A pharmaceutical composition in capsule form as claimed in Claim 7 in which said hydroxypropylmethylcellulose represents between about 15 to about 50 percent of the contents of said capsule.
- 9. A sustained-release pharmaceutical composition in capsule form as claimed in Claim 1 for oral administration of guanabenz, which comprises a pharmaceutically effective amount of guanabenz admixed with a hydroxypropylmethylcellulose wherein said hydroxypropylmethylcellulose has a methoxyl content of about 16-24 percent, a hydroxypropoxyl content of about 4-32 percent and an average number molecular weight above 50,000 and said hydroxypropylmethylcellulose represents between about 10 to about 60 weight percent of the contents of said capsule.

INTERNATIONAL SEARCH REPORT

	International Application No UI/UC	0070070			
I. CLASSIFICATION OF SUBJECT MATTER (If several classification)	cation symbols apply, indicate all) 3				
According to International Patent Classification (IPC) or to both Natio	onal Classification and IPC 0 4				
A61K 9/40, A61K 9/22. A61K 9/52 .					
II. FIELDS SEARCHED Minimum Document	tation Searched 4				
	Classification Symbols				
Classification System					
US 424/37, 19					
Documentation Searched other the to the Extent that such Documents	nan Minimum Documentation are Included in the Fields Searched 6				
III. DOCUMENTS CONSIDERED TO BE RELEVANT 14					
Category Citation of Document, 16 with Indication, where appr	ropriate, of the relevant passages 17	Relevant to Claim No. 18			
Y US,A, 3,065,143 to Christen 20 Nov. 1962		19			
y US,A, 3,870,790 to Lowey <u>et</u> 11 March 1975	al., published	1-9			
x,y US,A, 4,126,672 to Sheth <u>et</u> 21 Nov. 1978	al., published	1-9			
y US,A, 4,167,558 to Sheth <u>et</u> 11 Sept. 1979	al., published	1-9			
X,Y US,A, 4,173,626 to Dempski 6 Nov. 1979	et al., published	1-9			
y US,A, 4,226,849 to Schor, p 7 Oct. 1980	oublished	1-9			
Special categories of cited documents: 16 "A" document defining the general state of the art which is not	"T" later document published after or priority date and not in conficited to understand the princip	the international filing date lict with the application but le or theory underlying the			
"E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means	"X" document of particular relevat cannot be considered novel of involve an inventive step. "Y" document of particular relevat cannot be considered to involve document is combined with one ments, such combination being in the art.	nce; the claimed invention an inventive step when the e or more other such docu- obvious to a person skilled			
"P" document published prior to the international many date later than the priority date claimed	"&" document member of the same	potenti remity			
IV. CERTIFICATI N	1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	earch Report 9			
Date of the Actual Completion of the International Search ³	Date of Mailing of this International S 1 3 MAY 1985,	Bairii Isabais			
5 May 1985	Signafure of Authorized Officer 20-				
International Searching Authority 1	יייייייייייייייייייייייייייייייייייייי				

IS/US

વ

International Application PCT/US 8 5 / 00 3 0

III. DOCU	MENTS CONSIDERED T BE RELEVANT (CONTINUED FROM THE SEC NO SHEE	
Category •	Citation of Document, 16 with indication, where appropriate; of the relevant passages 17	Relevant to Claim No 18
Y	US,A, 4,389,393 to Schor et al., published 21 June 1983	1-9
P,X,Y	US,A, 4,478,819 to Hercelin et al., published 23 Oct. 1984	1-9
		•
•		
		1
į		
•		
	· ·	
		<u> </u>